

University of Groningen

Antibiotic resistance plasmids cointegrated into a megaplasmid harbouring the blaOXA-427 carbapenemase gene

Desmet, Stefanie; Nepal, Suruchi; van Dijl, Jan Maarten; Van Ranst, Marc; Chlebowicz, Monika A; Rossen, John W; Van Houdt, Jeroen K J; Maes, Piet; Lagrou, Katrien; Bathoorn, Erik

Published in:
Antimicrobial Agents and Chemotherapy

DOI:
[10.1128/AAC.01448-17](https://doi.org/10.1128/AAC.01448-17)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Desmet, S., Nepal, S., van Dijl, J. M., Van Ranst, M., Chlebowicz, M. A., Rossen, J. W., Van Houdt, J. K. J., Maes, P., Lagrou, K., & Bathoorn, E. (2018). Antibiotic resistance plasmids cointegrated into a megaplasmid harbouring the blaOXA-427 carbapenemase gene. *Antimicrobial Agents and Chemotherapy*, 62(3), [e01448-17]. <https://doi.org/10.1128/AAC.01448-17>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Antibiotic Resistance Plasmids Cointegrated into a Megaplasmid Harboring the *bla*_{OXA-427} Carbapenemase Gene

Stefanie Desmet,^a Suruchi Nepal,^b Jan Maarten van Dijk,^b Marc Van Ranst,^c Monika A. Chlebowicz,^b John W. Rossen,^b Jeroen K. J. Van Houdt,^d Piet Maes,^c Katrien Lagrou,^a Erik Bathoorn^b

^aDepartment of Microbiology and Immunology, Laboratory of Clinical Bacteriology and Mycology, KU Leuven-University of Leuven, Leuven, Belgium

^bUniversity of Groningen, University Medical Center Groningen, Department of Medical Microbiology, Groningen, The Netherlands

^cDepartment of Microbiology and Immunology, Laboratory of Clinical Bacteriology and Epidemiological Virology, KU Leuven-University of Leuven, Leuven, Belgium

^dCentre for Human Genetics, University Hospitals Leuven, KU Leuven, Leuven, Belgium

ABSTRACT OXA-427 is a new class D carbapenemase encountered in different species of *Enterobacteriaceae* in a Belgian hospital. To study the dispersal of this gene, we performed a comparative analysis of two plasmids containing the *bla*_{OXA-427} gene, isolated from a *Klebsiella pneumoniae* strain and an *Enterobacter cloacae* complex strain. The two IncA/C2 plasmids containing *bla*_{OXA-427} share the same backbone; in the *K. pneumoniae* strain, however, this plasmid is cointegrated into an IncFII plasmid, forming a 321-kb megaplasmid with multiple multiresistance regions.

KEYWORDS carbapenemase, megaplasmids, OXA-427

The dissemination of carbapenem resistance in *Enterobacteriaceae* is a major public health concern, as carbapenem antibiotics are a last-resort therapy for these common pathogens. The new OXA-427 carbapenemase presents a profile of wide resistance to β -lactams, including broad-spectrum penicillins, extended-spectrum cephalosporins, and carbapenems (1, 2). The *bla*_{OXA-427} gene was detected in different *Enterobacteriaceae* species ($n = 6$) and in different patients ($n = 9$) hospitalized at a university hospital in Belgium, in different hospital wards, between March 2012 and June 2014 (1); this suggests a high level of interspecies transmission of this resistance gene, probably associated with horizontal gene transfer of a specific plasmid. Recent studies, based on long-read sequencing data, suggest that mobility of resistance genes occurs at multiple nested genetic levels, with transmission of strains between individuals, transfer of plasmids between strains, and transposition of resistance genes between plasmids (3, 4). We report the comparative analysis of two plasmids containing the *bla*_{OXA-427} gene, for future study of the potential mechanism of interspecies transmission of the *bla*_{OXA-427} gene.

Two OXA-427-producing isolates, one of *Klebsiella pneumoniae* (KLPN57) and the other of the *Enterobacter cloacae* complex (ENCL58), were cultured from clinical samples from two different patients, in January 2013 and October 2013, respectively. Genomic DNA was extracted from these isolates and PacBio sequencing was performed at the Genomics Core laboratory (Leuven, Belgium), according to the protocol used by Conlan et al. (5). Genomic DNA was sheared into 10- to 15-kb fragments and converted into SMRTbell template libraries. Each strain was sequenced on a single SMRT cell with the PacBio RSII system, using P6 polymerase binding and C4 sequencing kits. The sequencing coverage for both plasmids ranged from 160-fold to 400-fold coverage. Genome assemblies were performed using HGAP 3 as part of SMRTAnalysis 2.3.0, annotation of the genes was performed by using RAST (<http://rast.nmpdr.org>), and the

Received 18 July 2017 Returned for modification 5 September 2017 Accepted 21 December 2017

Accepted manuscript posted online 8 January 2018

Citation Desmet S, Nepal S, van Dijk JM, Van Ranst M, Chlebowicz MA, Rossen JW, Van Houdt JKJ, Maes P, Lagrou K, Bathoorn E. 2018. Antibiotic resistance plasmids cointegrated into a megaplasmid harboring the *bla*_{OXA-427} carbapenemase gene. *Antimicrob Agents Chemother* 62:e01448-17. <https://doi.org/10.1128/AAC.01448-17>.

Copyright © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to Stefanie Desmet, stefanie.desmet@uzleuven.be.





FIG 1 Circular representation of pENCL58_01, harboring *bla*_{OXA-427}. Only genes for partitioning/replication (yellow), conjugative transfer (green), antibiotic resistance (red), and mobility (gray) are depicted. Prime symbols indicate truncated genes.

functional assignments for the predicted open reading frames (ORFs) were confirmed by using BLASTP with the Protein Data Bank (PDB) from NCBI (<http://www.ncbi.nlm.nih.gov>). IS finder and ResFinder were used for identification of mobile elements and antibiotic resistance genes, respectively (6, 7). PubMLST was used for multilocus sequence typing (MLST) (8). The plasmid maps were drawn by using SnapGene software.

The genomic content of *Enterobacter cloacae* strain ENCL58 consists of a 5.0-Mb genome with sequence type 171 (ST-171) and three plasmids (177 kb, 121 kb, and 84 kb). The 177,037-bp plasmid pENCL58_01, carrying the *bla*_{OXA-427} gene, is an IncA/C2 plasmid of plasmid multilocus sequence type 3, 100% identical to KX869741.1 (1) (Fig. 1). The other two plasmids (IncFII and IncFII types) carry no known resistance genes. The IncA/C2 plasmid shows overall high levels of homology with other IncA/C2 plasmids found in *Enterobacter aerogenes* (GenBank accession number [FO203354.1](#); 99% sequence identity and 87% coverage), *Klebsiella pneumoniae* (GenBank accession number [JQ010984.1](#); 99% sequence identity and 86% coverage), and *Salmonella en-*



terica (GenBank accession number [KM670336.1](#); 99% sequence identity and 75% coverage). The plasmid pENCL58_01 contains 223 coding sequences, including three functional antibiotic resistance genes (*bla*_{OXA-427}, *bla*_{TEM-1B'} and *strB*) and two truncated resistance genes (*strA* and *sul2*). The *bla*_{TEM-1} gene and the *strB-strA* gene combination are located adjacent to the *bla*_{OXA-427} resistance transposon, in the direction away from *repA*, and both are associated with the presence of IS15 elements, making this a resistance island (RI). In this plasmid, two transfer regions were identified, one located from position 49313 to position 57324 and including 3 *tra* genes and *kfrA* (for plasmid maintenance) and one located from position 145665 to position 173831 and including 12 *tra* genes and *trhF*.

aac.asm.org 3




FIG 3 Pairwise comparison of pENCL58_01 (upper sequence) and pKLPN57_01 (lower sequence). The *int1* gene of pENCL58_01 is used as a starting point. At the starting position, the homologous plasmid is disrupted and integrated into the pKLPN57_01 megaplasmid. Connecting bars represent regions with BLASTN matches of >99.9% in the same (red) and inverted (blue) orientations, visualized with the Artemis Comparison Tool (13).

The plasmid pKLPN57_01 contains resistance genes for aminoglycoside resistance (*aadA24*, *aacA4*, and *strB*), β -lactam resistance (*bla*_{OXA-10}, *bla*_{TEM-1B}, and *bla*_{OXA-427}), chloramphenicol resistance (*floR*), and sulfonamide resistance (*sul1*). Five additional resistance genes (*strA*, *bla*_{OXA-2}, *sul2*, *bla*_{OXA-10}, and *aacA4*) are truncated and potentially dysfunctional. Interestingly, two types of *repA* genes, coding for plasmid replication proteins, were detected on that plasmid. These *repA* genes belong to two distinct plasmid incompatibility groups, namely, IncA/C2 and IncFIIb. Further analysis confirmed that pKLPN57_01 is composed of a novel plasmid belonging to the IncFIIb group and an IncA/C2 plasmid highly similar to pENCL58_01, which was identified in *Enterobacter cloacae* ENCL58 (Fig. 3). Additionally, the plasmid carries two parts of a *repA* gene related to *colW* separated by the IncA/C2 region, together constituting the complete replication gene. Based on the presence of different *repA* genes and the large number of mobile genetic elements, the pKLPN57_01 plasmid is in fact a megaplasmid of 321 kb and could be a result of multiple fusions of different plasmids and/or mobile elements. The IncFIIb region showed the highest level of homology with a *K. pneumoniae* plasmid (GenBank accession number [CP021541.1](#); 99% identity and 68% coverage) and the *colW* region with a *Achromobacter xylosoxidans* subsp. *denitrificans* plasmid (GenBank accession number [HF679279.1](#); 100% identity and 37% coverage). Unfortunately, we were not able to identify the sources of these plasmid elements.

The genetic environment downstream of the *bla*_{OXA-427} gene, characterized by

IS1326 and a Tn3 family transposon, is identical in the two plasmids. Upstream of the *bla*_{OXA-427} gene, however, additional resistance genes are present in pKLPN57_01, compared to pENCL58_01 (Fig. 1 and 2).

In Fig. 3, the alignment of pENCL58 and pKLPN57 is shown. In the pKLPN57_01 megaplasmid, the *IncA/C2* region is flanked by identical class I integron integrase genes, *intI1*, at both sites. At the 5' site, the *intI1* gene is fused with a resistance integron. At the 3' site, the *intI1* gene of *IncA/C2* is fused with a partial *colW*-like *repA* gene by a transposase gene of the Tn3 family and an IS6 insertion sequence. In both regions of fusion, there is an abundance of Tn3 family transposases and IS6 and IS110 insertion sequences. The region homologous to pENCL58 is divided into two parts by a transposon flanked on both sides by IS91, carrying the resistance gene *floR*.

We conclude that the megaplasmid in the *K. pneumoniae* isolate is a result of cointegration of an *IncA/C2*-type plasmid harboring *bla*_{OXA-427} with an *IncF1b*-type plasmid. The ability of Tn3 and insertion sequences to mediate cointegration of plasmids is well known (9). Plasmid cointegrates have been reported previously for carbapenemase-producing *Enterobacteriaceae* strains (5, 10); they showed *IncFla*-type plasmid backgrounds and IS26 insertion sequences flanking the regions of fusion. To our knowledge, however, cointegrates formed from large resistance plasmids with sizes of >300 kb have not been reported previously. The plasmid cointegration seems to represent an important molecular pathway for interspecies transfer of plasmids and their resistance genes. Cointegration with disruption of replicons, as observed in pKLPN57_01, may allow plasmid incompatibility to be overcome (11). Next, cointegration with a plasmid that has coevolved in *K. pneumoniae* may provide the transferred plasmid with specific conjugation and restriction modification machinery (12). The cointegration of plasmids into pKLPN57_01 has probably taken place in recent years. Future studies will need to clarify whether megaplasmids of the size presented in our study are persistent or represent a temporary initial stage in the evolution of antibiotic resistance plasmids.

Accession number(s). The annotated plasmid sequences of pKLPN57_01 and pENCL58_01 were deposited in DDBJ/EMBL/GenBank under accession numbers [LT882698](#) and [LT882699](#), respectively.

ACKNOWLEDGMENTS

We thank Artur Sabat (University of Groningen, University Medical Center Groningen, Groningen, the Netherlands) for reviewing the sequence analysis and Y. Pfeifer (Robert Koch Institute) for kindly providing the *Escherichia coli* j53 azide-resistant strain.

REFERENCES

- Bogaerts P, Naas T, Saegeman V, Bonnin RA, Schuermans A, Evrard S, Bouchahrouf W, Jove T, Tande D, de Bolle X, Huang TD, Dortet L, Glupczynski Y. 2017. OXA-427, a new plasmid-borne carbapenem-hydrolysing class D β -lactamase in Enterobacteriaceae. *J Antimicrob Chemother* 72:2469–2477. <https://doi.org/10.1093/jac/dkx184>.
- Bogaerts P, Thierry N, Evrard S, Bouchahrouf W, Saegeman V, Lasserre C, Tande D, De Bolle X, Huang TD, Glupczynski Y. 2015. OXA-427, a new plasmidic ESBL class D OXA-carbapenemase recovered from Enterobacteriaceae clinical isolates. *Abstr ECCMID 2015*, abstr P1307.
- Sheppard AE, Stoesser N, Wilson DJ, Sebra R, Kasarskis A, Anson LW, Giess A, Pankhurst LJ, Vaughan A, Grim CJ, Cox HL, Yeh AJ, Sifri CD, Walker AS, Peto TE, Crook DW, Mathers AJ. 2016. Nested Russian doll-like genetic mobility drives rapid dissemination of the carbapenem resistance gene *bla*_{KPC}. *Antimicrob Agents Chemother* 60:3767–3778. <https://doi.org/10.1128/AAC.00464-16>.
- Mathers AJ, Stoesser N, Sheppard AE, Pankhurst L, Giess A, Yeh AJ, Didelot X, Turner SD, Sebra R, Kasarskis A, Peto T, Crook D, Sifri CD. 2015. *Klebsiella pneumoniae* carbapenemase (KPC)-producing *K. pneumoniae* at a single institution: insights into endemicity from whole-genome sequencing. *Antimicrob Agents Chemother* 59:1656–1663. <https://doi.org/10.1128/AAC.04292-14>.
- Conlan S, Thomas PJ, Deming C, Park M, Lau AF, Dekker JP, Snitkin ES, Clark TA, Luong K, Song Y, Tsai YC, Boitano M, Dayal J, Brooks SY, Schmidt B, Young AC, Thomas JW, Bouffard GG, Blakesley RW, Mullikin JC, Korlach J, Henderson DK, Frank KM, Palmore TN, Segre JA. 2014. Single-molecule sequencing to track plasmid diversity of hospital-associated carbapenemase-producing Enterobacteriaceae. *Sci Transl Med* 6:254ra126. <https://doi.org/10.1126/scitranslmed.3009845>.
- Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 34:D32–D36. <https://doi.org/10.1093/nar/gkj014>.
- Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
- Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Ponten T, Ussery DW, Aarestrup FM, Lund O. 2012.

- Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.
9. Shapiro JA. 1979. Molecular model for the transposition and replication of bacteriophage Mu and other transposable elements. *Proc Natl Acad Sci U S A* 76:1933–1937. <https://doi.org/10.1073/pnas.76.4.1933>.
 10. Chavda KD, Chen L, Jacobs MR, Rojzman AD, Bonomo RA, Kreiswirth BN. 2015. Complete sequence of a *bla*_{KPC}-harboring cointegrate plasmid isolated from *Escherichia coli*. *Antimicrob Agents Chemother* 59: 2956–2959. <https://doi.org/10.1128/AAC.00041-15>.
 11. Lin DL, Ramirez MS, Tran T, Tolmasky ME. 2013. A cointegrate-like plasmid that facilitates dissemination by conjugation of the extended-spectrum β -lactamase CTX-M-17. *Antimicrob Agents Chemother* 57: 5191–5192. <https://doi.org/10.1128/AAC.01365-13>.
 12. Ramirez MS, Traglia GM, Lin DL, Tran T, Tolmasky ME. 2014. Plasmid-mediated antibiotic resistance and virulence in Gram-negatives: the *Klebsiella pneumoniae* paradigm. *Microbiol Spectr* 2:1–15. <https://doi.org/10.1128/microbiolspec.PLAS-0016-2013>.
 13. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis Comparison Tool. *Bioinformatics* 21: 3422–3423. <https://doi.org/10.1093/bioinformatics/bti553>.